

GEOMETRICAL EVALUATION OF STAR PRODUCTS

Cosmas Zachos

High Energy Physics Division, Argonne National Laboratory, Argonne, IL 60439-4815, USA
zachos@hep.anl.gov

Abstract

The geometric picture of the star-product based on its Fourier representation kernel is utilized in the evaluation of chains of star-products and the intuitive appreciation of their associativity and symmetries. Such constructions appear even simpler for a variant asymmetric product, and carry through for the standard star-product supersymmetrization.

§1. Groenewold's noncommutative $\star$ -product [1] of phase-space functions $f(x, p)$ and $g(x, p)$ is the unique associative pseudodifferential deformation [2] of ordinary products:

$$\star \equiv e^{i\hbar(\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x)/2}. \quad (1)$$

It is the linchpin of deformation (phase-space) quantization [3, 2], as well as applications of matrix models and non-commutative geometry ideas in M-physics [4]. In practice, since it involves exponentials of derivative operators, it may be evaluated through translation of function arguments,

$$f(x, p) \star g(x, p) = f\left(x + \frac{i\hbar}{2} \overrightarrow{\partial}_p, p - \frac{i\hbar}{2} \overrightarrow{\partial}_x\right) g(x, p). \quad (2)$$


However, explicit evaluations of long strings of star-products in this language frequently appear intractable, unless the phase-space functions involved consist of exponentials or simple polynomials [5].

Baker [6] has utilized the more practical Fourier representation of this product as an integral kernel:

$$f \star g = \frac{1}{\hbar^2 \pi^2} \int dp' dp'' dx' dx'' f(x', p') g(x'', p'') \exp\left(\frac{-2i}{\hbar} (p(x' - x'') + p'(x'' - x) + p''(x - x'))\right). \quad (3)$$

The cyclic determinantal expression multiplying $-2i/\hbar$ in the exponent is twice the area of the phase-space triangle $(\mathbf{r}'', \mathbf{r}', \mathbf{r})$, where $\mathbf{r} \equiv (x, p)$, namely,

$$2A(\mathbf{r}'', \mathbf{r}', \mathbf{r}) = (\mathbf{r}' - \mathbf{r}) \wedge (\mathbf{r} - \mathbf{r}'') = \mathbf{r}'' \wedge \mathbf{r}' + \mathbf{r}' \wedge \mathbf{r} + \mathbf{r} \wedge \mathbf{r}''. \quad (4)$$



In this representation, multiple star-products turn out to be simpler to evaluate, and the geometrical constructions they motivate exhibit conspicuously the symmetries and the associativity of these products.

The representation thus rises to the level of a ‘picture’, in Dirac’s sense of a “way of looking at the fundamental laws which makes their self-consistency obvious” [7]. Such evaluations are illustrated below, with some practical hints, for the standard star-product, as well as for some common variants and extensions.


§2. In the Fourier representation, a triple star-product can be expressed relatively simply,

$$f \star g \star h = \frac{1}{\hbar^4 \pi^4} \int d\bar{p} dp' dp'' dp''' d\bar{x} dx' dx'' dx''' f(x', p') g(x'', p'') h(x''', p''') \exp \frac{-4i}{\hbar} (A(r'', r', \bar{r}) + A(r''', \bar{r}, r)) , \quad (5)$$

while the intermediary $d\bar{x} d\bar{p}$ integrations collapse to δ -functions:

$$\begin{aligned} f \star g \star h &= \frac{1}{\hbar^2 \pi^2} \int dp' dp'' dp''' dx' dx'' dx''' f(x', p') g(x'', p'') h(x''', p''') \times \\ &\times \delta(x - x' + x'' - x''') \delta(p - p' + p'' - p''') \exp \left(\frac{-4i}{\hbar} A(r''', r'', r') \right) . \end{aligned} \quad (6)$$

The product thus amounts to a triangle and a point for the effective phase-space argument $\mathbf{r} = (x, p)$, which lies on the new vertex of the parallelogram resulting from doubling up the triangle $(\mathbf{r}''', \mathbf{r}'', \mathbf{r}')$, such that $\mathbf{r}' - \mathbf{r}'''$ is one diagonal. The effective argument $\mathbf{r}$ lies at the end of the *other* diagonal, starting from $\mathbf{r}''$,



It is then straightforward to note how this expression bears no memory of the grouping (order of association) in which the two star-multiplications were performed, since the vertex $\mathbf{r}$ of the parallelogram is reached from $\mathbf{r}'''$ by translating through $\mathbf{r}' - \mathbf{r}''$, or, equivalently, from $\mathbf{r}'$ by translating through $\mathbf{r}''' - \mathbf{r}''$. As a result, this may well realize the briefest graphic proof of the distinctive associativity property of the star-product,

$$(f \star g) \star h = f \star (g \star h) . \quad (7)$$


The symmetries of the triple star-product, (1-3 complex conjugacy, effective cyclicity, e.t.c.) are now evident by inspection. Moreover, integration of this triple product with respect to the effective argument (x, p) (tracing), e.g. to yield a lagrangian interaction term, trivially eliminates the δ -function to result in a compact cyclic expression of the above triangle construction for the three functions star-multiplied,

$$\int dx dp f \star g \star h = \frac{1}{\hbar^2 \pi^2} \int dp_1 dp_2 dp_3 dx_1 dx_2 dx_3 f(x_1, p_1) g(x_2, p_2) h(x_3, p_3) \exp \left(\frac{-4i}{\hbar} A(r_3, r_2, r_1) \right) . \quad (8)$$

A four function star-product (with three stars) involves the sum of the areas of two triangles, $(\mathbf{r}_3, \mathbf{r}_2, \mathbf{r}_1)$ and $(\mathbf{r}, \mathbf{r}_4, \mathbf{r}_1 - \mathbf{r}_2 + \mathbf{r}_3)$. A five function star-product involves the exponential of the sum of areas of two triangles, $(\mathbf{r}_3, \mathbf{r}_2, \mathbf{r}_1)$ and $(\mathbf{r}_5, \mathbf{r}_4, \mathbf{r}_1 - \mathbf{r}_2 + \mathbf{r}_3)$, with the effective argument restricted by $\delta(\mathbf{r} - \mathbf{r}_1 + \mathbf{r}_2 - \mathbf{r}_3 + \mathbf{r}_4 - \mathbf{r}_5) \equiv \delta(\mathbf{r} - \mathbf{s}_5)$. Recursively, so on for even numbers of star-multiplied functions, the phase involving the sums $A(\mathbf{r}_3, \mathbf{r}_2, \mathbf{r}_1) + A(\mathbf{r}_5, \mathbf{r}_4, \mathbf{s}_3) + \dots + A(\mathbf{r}, \mathbf{r}_{2n}, \mathbf{s}_{2n-1})$. Respectively, for odd numbers of functions,

the phase involving sums $A(\mathbf{r}_3, \mathbf{r}_2, \mathbf{r}_1) + A(\mathbf{r}_5, \mathbf{r}_4, \mathbf{s}_3) + \dots + A(\mathbf{r}_{2n+1}, \mathbf{r}_{2n}, \mathbf{s}_{2n-1})$, with effective phase-space argument restrictions to $\mathbf{r} = \mathbf{s}_{2n+1} \equiv \sum_{m=1}^{2n+1} (-)^{m+1} \mathbf{r}_m$.

As an illustration, consider phase-space points $\mathbf{r}_i$ arrayed in a regular zigzag pattern, (i.e. for the star-multiplied functions getting support only on those points on the zigzag). The arguments of the δ -functions, $\mathbf{s}_{2n+1}$, then lie on a line, while the areas of the triangles demarcated by these points increase in regular arithmetic progression ($A, 2A, 3A, 4A, \dots$):



This result is independent of the pitch of the zigzag, i.e. the angle at $\mathbf{r}_2$ —which, in this figure, is chosen to be $\pi/2$, since this is a local maximum of the areas A of the triangles for variable pitch but fixed lengths $\mathbf{r}_i - \mathbf{r}_{i+1}$. One might well wonder if the configuration pictured could be used to define a “classical path”: its contribution to the phase of the exponential through the sum of all triangle areas, $(1 + 2 + 3 + 4 + \dots)A$, is stationary with respect to variations such as this angle variation discussed. The question suggests itself, then, whether configurations stationary under *all* variations can be constructed, leading to a stationary phase evaluation of large/infinite star-products, e.g. useful in evaluating star-exponentials (which yield time-evolution operators in phase-space [2]); but, so far, no cogent general answers appear at hand.

§3. A variant of the star-product (cohomologically equivalent to it) is the lopsided associative product of Voros [8],

$$\bowtie \equiv e^{i\hbar \overleftarrow{\partial}_x \overrightarrow{\partial}_p} . \quad (9)$$

It is sometimes convenient to rotate phase-space variables canonically (i.e. preserving their Poisson Brackets),

$$(x, p) \mapsto \left(\frac{x + ip}{\sqrt{-2i}}, \frac{x - ip}{\sqrt{-2i}} \right) , \quad (10)$$

to represent this product as

$$\begin{aligned} \bowtie &\equiv e^{\hbar(\overleftarrow{\partial}_x - i\overleftarrow{\partial}_p)(\overrightarrow{\partial}_x + i\overrightarrow{\partial}_p)/2} = e^{i\hbar(\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x)/2} e^{i\hbar(\overleftarrow{\partial}_x \overrightarrow{\partial}_x + \overleftarrow{\partial}_p \overrightarrow{\partial}_p)/2} \\ &= \star e^{-\hbar(\overleftarrow{\partial}_x^2 + \overleftarrow{\partial}_p^2)/4} e^{-\hbar(\overrightarrow{\partial}_x^2 + \overrightarrow{\partial}_p^2)/4} e^{\hbar((\overleftarrow{\partial}_x + \overrightarrow{\partial}_x)^2 + (\overleftarrow{\partial}_p + \overrightarrow{\partial}_p)^2)/4} . \end{aligned} \quad (11)$$

This turns out to be the covariant transform of the $\star$ -product which controls the dynamics when Wigner distributions are transformed into Husimi distributions [9], a smoothed representation popular in applications. It is plain that the Gaussian-Laplacian factors $T^{-1}(\partial_x, \partial_p) \equiv \exp(-\hbar(\partial_x^2 + \partial_p^2)/4)$ merely dress the standard star-product into Voros' product [8],


$$T(f \star g) = T(f) \bowtie T(g) . \quad (12)$$

Consequently, the Lie algebra of brackets of $\bowtie$ is isomorphic to the Moyal algebra [3] (the algebra of brackets of $\star$, i.e. $\{\{f, g\}\} \equiv f \star g - g \star f$), in comportance with the general result on the essential uniqueness of the Moyal algebra as the one-parameter deformation of the Poisson Bracket algebra [10].

Actually, in Fourier space, this product in its original representation (9) appears simpler than the $\star$ -product,

$$f \bowtie g = \frac{1}{2\pi\hbar} \int d\mathbf{r}' d\mathbf{r}'' f(x', p') g(x'', p'') \delta(x'' - x) \delta(p' - p) \exp\left(\frac{i}{\hbar}(x'' - x')(p' - p'')\right) . \quad (13)$$


The phase-space integral is then effectively a two-dimensional $\int dx' dp''$, not a four-dimensional one, as the kernel has vanishing support everywhere but on the lines $x'' = x, p' = p$. The triangle whose doubled area multiplies $-i/\hbar$ in the exponent is now a phase-space *right* triangle $(\mathbf{r}'', \mathbf{r}', \mathbf{r})$, with its side $\mathbf{r} - \mathbf{r}'$ horizontal, and its side $\mathbf{r} - \mathbf{r}''$ vertical:



The triple product is then seen to be actuating shifts on a rectangular lattice,

$$\begin{aligned} (f \bowtie g) \bowtie h &= \frac{1}{(2\pi\hbar)^2} \int d\mathbf{r}' d\mathbf{r}'' d\mathbf{r}''' f(x', p') g(x'', p'') h(x''', p''') \times \\ &\times \delta(x''' - x) \delta(p' - p) \exp\left(\frac{i}{\hbar}(x'(p'' - p') + x''(p''' - p'') + x'''(p' - p'''))\right) . \end{aligned} \quad (14)$$

The phase is a cyclic expression with no memory of the order of association, which thus proves associativity for this product, $(f \bowtie g) \bowtie h = f \bowtie (g \bowtie h)$. Pictorially, the phase is the area of the entire encompassing rectangle with diagonal $\mathbf{r}''' - \mathbf{s}$, minus the area of the rectangle with diagonal $\mathbf{r}' - \mathbf{r}''$; which is also equal to the *sum* of the areas of the rectangles with diagonals $\mathbf{s}' - \mathbf{r}'$, and $\mathbf{r}''' - \mathbf{r}''$, respectively. (In general, it is not twice the area of the triangle $(\mathbf{r}', \mathbf{r}'', \mathbf{r}''')$.)



The construction for an n -tuple $\bowtie$ -product follows simply,

$$\frac{1}{(2\pi\hbar)^n} \int d\mathbf{r}_1 \dots d\mathbf{r}_n f_1(r_1) \dots f_n(r_n) \delta(x_n - x) \delta(p_1 - p) \exp \left(\frac{i}{\hbar} \sum_{m=1}^n x_m (p_{m+1} - p_m) \right), \quad (15)$$

where p_{n+1} is defined as p_1 . One may note the effective nearest-neighbor interaction in the chain suggested.

§4. A superspace generalization of the star-product was introduced in ref [11], (to codify the graded extension of Moyal's algebra introduced in ref [12]),

$$(1 + \hbar \overleftarrow{\partial}_\theta \overrightarrow{\partial}_\theta) \star \equiv \diamond \star. \quad (16)$$

Here, θ is the superspace Grassmann variable (nilpotent, and commuting with the phase-space variables): the extended star-product is then a direct product of the conventional piece with a superspace factor $1 + \hbar \overleftarrow{\partial}_\theta \overrightarrow{\partial}_\theta$. Thus, the above extended product could have been alternatively written as

$$e^{\hbar \overleftarrow{\partial}_\theta \overrightarrow{\partial}_\theta} \star. \quad (17)$$

Hence, it can also be rewritten [13] as the evocative form,

$$e^{\frac{i\hbar}{2} (\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x) + \hbar \overleftarrow{\partial}_\theta \overrightarrow{\partial}_\theta}. \quad (18)$$

Nevertheless, the original form displays associativity more readily, since the factor acting on the Grassmann structure is patently associative,

$$(A \diamond B) \diamond C = A \diamond (B \diamond C), \quad (19)$$

acting on 1d bosonic superfields $A(\theta) = a + \theta\alpha$, $B(\theta) = b + \theta\beta$, so that

$$A \diamond B = ab + \hbar\alpha\beta + \theta(\alpha b + a\beta). \quad (20)$$

Note the loose analogy to complex multiplication $\bar{z}_1 z_2$. Even though this analogy cannot rise to an isomorphism, as evident from its noncommutativity and longer products such as the above, still, it turns out to be useful for actual evaluation of products in collecting the Grassmann even and odd terms in the answer. The symmetry of this product is further displayed by setting $\hbar = 1$ and considering standard Grassmann Fourier transforms from bosonic to fermionic superfields, $\tilde{A}(\theta) = \int d\phi (1 + \phi\theta) A(\phi) = \alpha + \theta a$:

$$A \diamond B = \tilde{A} \diamond \tilde{B}. \quad (21)$$

Acknowledgments

Appreciation of discussions with T Curtright and D Fairlie is recorded. This work was supported in part by the US Department of Energy, Division of High Energy Physics, Contract W-31-109-ENG-38.

References

- [1] H Groenewold, *Physica* **12** (1946) 405-460
- [2] Further explored by F Bayen, M Flato, C Fronsdal, A Lichnerowicz, and D Sternheimer, *Ann Phys* **111** (1978) 61-110; F Hansen, *Rept Math Phys* **19** (1984) 361-381; reviewed by C Roger and V Ovsienko, *Russ Math Surv* **47** (1992) 135-191
- [3] J Moyal, *Proc Camb Phil Soc* **45** (1949) 99-124
- [4] N Seiberg and E Witten, *JHEP* **9909** (1999) 032
- [5] T Curtright, D Fairlie, and C Zachos, *Phys Rev* **D58** (1998) 025002
- [6] G Baker, *Phys Rev* **109** (1958) 2198-2206; formally underlain by J v Neumann, *Math Ann* **104** (1931) 570-578; also see D Fairlie, *Proc Camb Phil Soc* **60** (1964) 581-586; further see review by F Berezin, *Sov Phys Usp* **23** (1980) 763-787
- [7] P A M Dirac, *The Principles of Quantum Mechanics*, 4th edition, (Oxford University Press, 1958), Ch I.4
- [8] F Bayen, in *Group Theoretical Methods in Physics*, W Beiglböck et al (eds), *Lecture Notes in Physics* **94**, (Springer-Verlag, Heidelberg, 1979), pp 260-271; A Voros, *Phys Rev* **A40** (1989) 6814-6825
- [9] K Takahashi, *Prog Theo Phys Suppl* **98** (1989) 109-156
- [10] J Vey, *Comment Math Helv* **50** (1975) 421-454; W Arveson, *Comm Math Phys* **89** (1983) 77-102; P Fletcher, *Phys Lett* **B248** (1990) 323-328
- [11] D Fairlie and C Zachos, *Phys Lett* **B224** (1989) 101-107
- [12] D Fairlie, P Fletcher, and C Zachos, *Phys Lett* **B218** (1989) 203-206
- [13] E Fradkin and V Linetsky, *Mod Phys Lett* **A6** (1991) 217-224; also see F Berezin and M Marinov, *Ann Phys* **104** (1977) 336-362; I Batalin and E Fradkin, *Riv Nuovo Cim* **9** n10 (1986) 1-48